

# Equivalence of pointwise and global ellipticity estimates

Hendrik Vogt

## Abstract

Defining an elliptic operator  $-\nabla \cdot (a\nabla)$  via the form method one normally imposes pointwise conditions on the matrix valued function  $a$  in order to get positivity, ellipticity and sectoriality of the form. In this note we show that the pointwise conditions on  $a$  are equivalent to the corresponding global ones on the form.

MSC 1991: 35J20 (primary)

Let  $\Omega$  be an arbitrary open subset of  $\mathbb{R}^d$ ,  $a: \Omega \rightarrow \mathbb{C}^{d \times d}$  a locally integrable, hermitian matrix valued function. Define the symmetric form  $\tau$  in  $L_2(\Omega)$  by  $\tau(u) := \int a \nabla u \cdot \nabla \bar{u}$  on  $D(\tau) := C_c^\infty(\Omega)$ . If  $\tau$  is positive and closable then, by the form representation theorem,  $\bar{\tau}$  is associated with a positive selfadjoint operator in  $L_2(\Omega)$  (which corresponds to Dirichlet boundary conditions). The main aim of this note is to show that the positivity of the form  $\tau$  is equivalent to the positivity of the function  $a$ , i.e.,  $a \geq 0$  in the matrix sense a.e.

A case of particular interest is the following: Let  $a_1: \Omega \rightarrow \mathbb{R}^{d \times d}$  be locally integrable, symmetric matrix valued and locally strictly elliptic, i.e., for every compact set  $K \subseteq \Omega$  there exists  $\sigma > 0$  such that  $a_1(x) \geq \sigma$  in the matrix sense for almost all  $x \in K$ . Then it is known that

$$D(\tau_N) := \{u \in L_2(\Omega) \cap W_{2,loc}^1(\Omega); \tau_N(u) := \int a_1 \nabla u \cdot \nabla \bar{u} < \infty\}$$

defines a symmetric Dirichlet form in  $L_2(\Omega)$  (cf. [1, Thm. 1.3.9]; one can show the closedness of  $\tau_N$  like the completeness of the Sobolev space  $W_2^1(\Omega)$  because of the local strict ellipticity of  $a_1$ ). The associated selfadjoint operator in  $L_2(\Omega)$  corresponds to Neumann boundary conditions.

*Notation.* Let  $|M|$  denote the Lebesgue measure of a measurable set  $M \subseteq \mathbb{R}^d$ ,  $\chi_M$  the characteristic function of  $M$ .  $S_{d-1}$  is the unit sphere of  $\mathbb{R}^d$ , for the spectral radius of a hermitian matrix  $A \in \mathbb{C}^{d \times d}$  we write  $|A| (= \sup_{\xi \in S_{d-1}} |A\xi \cdot \xi|)$ . For a function  $f: \Omega \rightarrow \mathbb{R}$  we use the shorthand  $[f > 0]$  for the set  $\{x \in \Omega; f(x) > 0\}$  (and similarly  $[f < g]$  etc).  $Q(f)$  denotes the form domain of the multiplication operator  $f$  in  $L_2(\Omega)$ .

**Theorem.** Let  $a: \Omega \rightarrow \mathbb{C}^{d \times d}$  be a locally integrable hermitian matrix valued function,  $\tau(u) := \int a \nabla u \cdot \nabla \bar{u}$  for  $u \in C_c^\infty(\Omega)$ .

(a) The following are equivalent:

- (i)  $a \geq 0$  a.e.,
- (ii)  $\tau(u) \geq 0$  for all  $u \in C_c^\infty(\Omega)$ ,
- (iii)  $\tau$  is bounded from below, i.e., there exists  $c > 0$  such that  $\tau(u) \geq -c\|u\|_2^2$  for all  $u \in C_c^\infty(\Omega)$ .

(b) Let  $D$  be a sublattice of  $W_{1,loc}^1(\Omega) \cap L_2(\Omega)$  with  $C_c^\infty(\Omega) \subseteq D \subseteq \{u; a \nabla u \cdot \nabla \bar{u} \in L_1(\Omega)\}$ . Let  $V: \Omega \rightarrow [0, \infty)$  be a measurable function such that  $D \cap Q(V)$  is dense in  $L_2(\Omega)$ . Define the form  $\tau_0$  in  $L_2(\Omega)$  by  $\tau_0(u) := \int a \nabla u \cdot \nabla \bar{u} + \int V|u|^2$  on  $D(\tau_0) := D \cap Q(V)$ . Then  $\tau_0$  is positive if and only if  $a \geq 0$  in the matrix sense a.e.

**Remark.** Let  $a_1$  and  $\tau_N$  be as in the introduction. Assume that  $a_1 \xi \cdot \xi \geq |a \xi \cdot \xi|$  a.e. for all  $\xi \in \mathbb{R}^d$  (e.g.  $a_1 = (|a| + 1)I$ ). Then  $D := D(\tau_N)$  is an example of a sublattice  $D$  satisfying the assumption of part (b) of the theorem (or any sublattice  $D$  of  $D(\tau_N)$  with  $C_c^\infty(\Omega) \subseteq D$ ). A particular example is  $D = W_{\infty,c}^1(\Omega)$ , the space of Lipschitz continuous functions with compact support.

**Proof of the Theorem.** The only nontrivial part of (a) is showing that (iii) implies (i). For this purpose we assume that  $[a \not\geq 0]$  has positive measure. In the following we construct a function violating (iii), later on we modify this function in order to show that the form  $\tau_0$  defined in (b) is not positive.

Let  $(\xi_k)$  be a dense sequence in  $S_{d-1}$ . If  $a(x) \not\geq 0$  for some  $x \in \Omega$  then there exists  $k \geq |a(x)|$  such that  $a(x)\xi_k \cdot \xi_k \leq -1/k$ . We obtain

$$[a \not\geq 0] = \bigcup_{k \in \mathbb{N}} ([a\xi_k \cdot \xi_k \leq -1/k] \cap [|a| \leq k]),$$

so we can fix  $k \in \mathbb{N}$  such that  $F := [a\xi_k \cdot \xi_k \leq -1/k] \cap [|a| \leq k]$  has positive measure.

Now we localize  $F$ : Almost all  $x \in \Omega$  are Lebesgue points of the function  $\chi_F$ , so since  $F$  has positive measure there exists  $x \in \Omega$  satisfying

$$\frac{1}{|B_\delta(x)|} \int_{B_\delta(x)} \chi_F \rightarrow \chi_F(x) = 1 \quad \text{for } \delta \rightarrow 0.$$

Let  $\varepsilon \in (0, 1)$  be given. Fix  $\delta > 0$  in such a way that  $B := B_\delta(x) \subseteq \overline{B_\delta(x)} \subseteq \Omega$  and  $|B \cap F| = \int_B \chi_F \geq (1 - \varepsilon)|B|$ . Since the function  $|a|$  restricted to  $B$  is integrable, there exists  $\alpha > 0$  such that for all measurable sets  $M \subseteq B$  with  $|M| \leq \alpha$  we have  $\int_M |a| \leq \varepsilon|B|$ . By the regularity of the Lebesgue measure there exists an open set  $\Omega_0$ ,  $B \cap F \subseteq \Omega_0 \subseteq B$ , with  $|\Omega_0 \setminus F| \leq \alpha$ . It follows that

$$\int_{\Omega_0 \setminus F} |a| \leq \varepsilon|B| \quad \text{and} \quad |\Omega_0| \geq |B \cap F| \geq (1 - \varepsilon)|B|. \quad (1)$$

We can choose functions  $u_n \in C_c^\infty(\Omega_0)$  satisfying

$$0 \leq u_n \leq \frac{1}{n}, \quad \|\nabla u_n\|_\infty \leq 1, \quad \text{and} \quad |[\nabla u_n \neq \pm \xi_k] \cap \Omega_0| \leq \varepsilon |B|$$

for all  $n \in \mathbb{N}$ . (The reader should think of pieces of hyperplanes orthogonal to  $\xi_k$  which have distance  $\frac{2}{n}$  to each other and to the boundary of  $\Omega_0$ . On these hyperplanes set  $u_n := \frac{1}{n}$  and extend like a wash board.) The important point is that on the set  $G := F \cap [\nabla u_n = \pm \xi_k]$  the function  $a \nabla u_n \cdot \nabla u_n$  is less or equal  $-1/k$ .

Let  $H := \Omega_0 \setminus G$ . Then  $|H| \leq |\Omega_0 \setminus F| + |\Omega_0 \cap [\nabla u_n \neq \pm \xi_k]| \leq 2\varepsilon |B|$  since  $|\Omega_0 \setminus F| \leq |B \setminus F| \leq \varepsilon |B|$  by (1). We can estimate

$$\tau(u_n) = \int_G a \xi_k \cdot \xi_k + \int_H a \nabla u_n \cdot \nabla u_n \leq -\frac{1}{k} |G| + \int_{H \cap F} |a| + \int_{H \setminus F} |a|.$$

By (1) we obtain  $|G| = |\Omega_0| - |H| \geq (1 - 3\varepsilon) |B|$  and (since  $|a| \leq k$  on  $F$ )

$$\tau(u_n) \leq -\frac{1}{k} |G| + k |H \cap F| + \int_{\Omega_0 \setminus F} |a| \leq \left(-\frac{1}{k}(1 - 3\varepsilon) + 2k\varepsilon + \varepsilon\right) |B|. \quad (2)$$

Note that  $\|u_n\|_2 \leq \frac{1}{n} \|\chi_B\|_2$  since  $0 \leq u_n \leq \frac{1}{n}$ . Therefore, given  $c > 0$ , it is easy to choose first  $\varepsilon$  and then  $n$  in such a way that  $\tau(u_n) + c\|u_n\|_2^2 < 0$ . This completes the proof of (a).

Now we use the sequence  $(u_n)$  in order to construct a function  $u \in D(\tau_0)$  with  $\tau_0(u) < 0$ , thus proving (b). Let  $\varepsilon := 1/(2k^2 + 5k + 3)$ . Since  $D(\tau_0)$  is a dense sublattice of  $L_2(\Omega)$ , there exists a function  $0 \leq \varphi \in D(\tau_0)$  satisfying  $|\{\varphi < \chi_B\}| < \alpha$ . Hence

$$\int_{[\varphi < \chi_B]} |a| \leq \varepsilon |B| \quad (3)$$

according to the choice of  $\alpha$ . Since  $\varphi \in Q(V)$  and  $0 \leq u_n \leq \frac{1}{n}$  ( $n \in \mathbb{N}$ ) there exists  $n_0 \in \mathbb{N}$  with

$$\int V |\varphi \wedge u_n|^2 < \varepsilon |B| \quad \text{for all } n \geq n_0. \quad (4)$$

By the assumptions on  $D$  we have  $a \nabla \varphi \cdot \nabla \varphi \in L_1(\Omega)$ , so there exists  $\alpha_2 > 0$  such that for all measurable sets  $M \subseteq \Omega$  with  $|M| \leq \alpha_2$  we have  $\int_M |a \nabla \varphi \cdot \nabla \varphi| \leq \varepsilon |B|$ . Obviously  $\bigcap_{n \in \mathbb{N}} [0 < \varphi < 1/n] = \emptyset$ , so since  $u_n \leq 1/n$  there exists  $n \geq n_0$  with  $|[0 < \varphi < u_n]| \leq \alpha_2$ . Noting  $\nabla \varphi = 0$  a.e. on  $[\varphi = 0]$ , we obtain

$$\int_{[\varphi < u_n]} |a \nabla \varphi \cdot \nabla \varphi| = \int_{[0 < \varphi < u_n]} |a \nabla \varphi \cdot \nabla \varphi| \leq \varepsilon |B|. \quad (5)$$

For  $u := \varphi \wedge u_n \in D(\tau_0)$  we estimate (noting  $\nabla u = \nabla u_n$  a.e. on  $[u = u_n]$  and  $\nabla u = \nabla \varphi$  a.e. on  $[u < u_n] = [\varphi < u_n]$ )

$$\begin{aligned} \tau(u) &= \int_{[u=u_n]} a \nabla u \cdot \nabla u + \int_{[u < u_n]} a \nabla u \cdot \nabla u \\ &\leq \int a \nabla u_n \cdot \nabla u_n + \int_{[\varphi < u_n]} |a \nabla u_n \cdot \nabla u_n| + \int_{[\varphi < u_n]} |a \nabla \varphi \cdot \nabla \varphi|. \end{aligned}$$

By (3) and (5) we conclude  $\tau(u) \leq \tau(u_n) + 2\varepsilon|B|$ . From (4) recall  $\int V|u|^2 \leq \varepsilon|B|$ . According to (2) and the choice of  $\varepsilon$  it follows that

$$\tau_0(u) \leq \tau(u_n) + 3\varepsilon|B| \leq -\varepsilon|B| < 0.$$

This completes the proof of (b).  $\square$

Now let  $a: \Omega \rightarrow \mathbb{C}^{d \times d}$  be a locally integrable matrix valued function (no longer assumed to be hermitian). Define the sesquilinear form  $\tau_a$  in  $L_2(\Omega)$  by  $\tau_a(u) := \int a \nabla u \cdot \nabla \bar{u}$  on  $C_c^\infty(\Omega)$ . We call the function  $a$  *sectorial* if  $|\operatorname{Im}(a\xi \cdot \bar{\xi})| \leq \alpha \operatorname{Re}(a\xi \cdot \bar{\xi})$  a.e. for all  $\xi \in \mathbb{C}^d$  and some  $\alpha \geq 0$  and *strictly elliptic* if  $\operatorname{Re}(a\xi \cdot \bar{\xi}) \geq \sigma|\xi|^2$  a.e. for all  $\xi \in \mathbb{C}^d$  and some  $\sigma > 0$ . The form  $\tau_a$  is called *sectorial* if  $|\operatorname{Im} \tau_a(u)| \leq \alpha \operatorname{Re} \tau_a(u) + c\|u\|_2^2$  and *strictly elliptic* if  $\operatorname{Re} \tau_a(u) \geq \sigma\|\nabla u\|_2^2 - c\|u\|_2^2$  for all  $u \in C_c^\infty(\Omega)$  and some  $c \in \mathbb{R}$ ,  $\alpha \geq 0$ ,  $\sigma > 0$ .

**Corollary.** *The form  $\tau_a$  is sectorial (strictly elliptic) if and only if  $a$  is sectorial (strictly elliptic).*

*Proof.* Let  $a(x)^*$  denote the adjoint matrix of  $a(x)$ . Then  $a_R := (a + a^*)/2$  and  $a_I := (a - a^*)/2i$  are hermitian matrix valued functions. One calculates

$$\operatorname{Re}(a\xi \cdot \bar{\xi}) = a_R \xi \cdot \bar{\xi} \quad \text{and} \quad \operatorname{Im}(a\xi \cdot \bar{\xi}) = a_I \xi \cdot \bar{\xi} \quad (6)$$

for all  $\xi \in \mathbb{C}^d$ , so  $\operatorname{Re} \tau_a = \tau_{a_R}$  and  $\operatorname{Im} \tau_a = \tau_{a_I}$ . The form  $\tau_a$  is sectorial with  $\alpha$  as above if and only if the forms  $\alpha \operatorname{Re} \tau_a \pm \operatorname{Im} \tau_a = \tau_{\alpha a_R \pm a_I}$  are bounded from below. By the above theorem this is equivalent to  $\alpha a_R \pm a_I \geq 0$  a.e., i.e. to the sectoriality of  $a$  according to (6).

For the second part of the Corollary just apply the theorem to the function  $a_R - \sigma$ .  $\square$

**Acknowledgements.** The research of the author was supported by the Deutsche Forschungsgemeinschaft. The support is gratefully acknowledged. The author would like to thank Jürgen Voigt for a number of valuable remarks.

## References

- [1] E. B. DAVIES, *Heat kernels and spectral theory*, Cambridge Univ. Press, 1989.

*Fachrichtung Mathematik  
Institut für Analysis  
Technische Universität Dresden  
D-01062 Dresden  
Germany  
vogt@math.tu-dresden.de*